

Flat and étale morphisms

Maxim Mornev

All rings are commutative.

Contents

1 Flat morphisms	1
1.1 Preliminaries on tensor product	1
1.2 Flat modules	2
1.3 Artin-Rees lemma and Krull intersection theorem	4
1.4 Modules of finite length	5
1.5 Criteria of flatness	6
1.6 Flatness in the context of schemes	8
2 Étale morphisms	9
2.1 The module of Kähler differentials	9
2.2 Étale algebras over fields	13
2.3 Unramified morphisms	15
2.4 Étale morphisms	16

1 Flat morphisms

1.1 Preliminaries on tensor product

Let A be a ring, M and A -module. For all A -modules N_1, N_2 we have a natural isomorphism

$$\mathrm{Hom}_A(N_1 \otimes_A M, N_2) \cong \mathrm{Hom}_A(N_1, \mathrm{Hom}_A(M, N_2)).$$

In other words $\otimes_A M$ is left adjoint to $\mathrm{Hom}_A(M, -)$. Hence $\otimes_A M$ is right exact and commutes with colimits.

Left derived functors $L^i(\otimes_A M)(-)$ are denoted $\text{Tor}_i^A(-, M)$. A morphism of modules $M \rightarrow M'$ induces natural morphisms $\text{Tor}_i^A(-, M) \rightarrow \text{Tor}_i^A(-, M')$, so Tor_i is a bifunctor. The most important property of Tor is its commutativity:

Theorem 1.1.1. *Let A be a ring, and let M, N be A -modules. For every $i \geq 0$ there exists a natural isomorphism $\text{Tor}_i^A(N, M) \rightarrow \text{Tor}_i^A(M, N)$.¹*

We will not need the full force of this theorem and so omit its proof.

Proposition 1.1.2. *Let A be a ring, $I \subset A$ an ideal, and M an A -module. $\text{Tor}_1^A(A/I, M) = \ker(I \otimes_A M \rightarrow M)$.*

Proof. The short exact sequence $0 \rightarrow I \rightarrow A \rightarrow A/I \rightarrow 0$ induces an exact sequence $0 = \text{Tor}_1^A(A, M) \rightarrow \text{Tor}_1^A(A/I, M) \rightarrow I \otimes_A M \rightarrow M$. \square

Corollary 1.1.3. *Let $a \in A$ be a nonzero element. $\text{Tor}_1^A(A/(a), M)$ is the a -torsion of M .*

Let A, B be rings, N_1 an A -module, N_2 an A, B -bimodule, and N_3 a B -module. There is an isomorphism of A, B -bimodules

$$(N_1 \otimes_A N_2) \otimes_B N_3 \rightarrow N_1 \otimes_A (N_2 \otimes_B N_3),$$

which is natural in N_1, N_2, N_3 .

Also recall that if A is a ring and $S \subset A$ a multiplicative system, then the functor $\otimes_A A_S$ is isomorphic to the functor of localization at S .

1.2 Flat modules

Definition 1.2.1. Let A be a ring. A module M over A is called flat if $\otimes_A M$ is exact.

Proposition 1.2.2. *Let $A \rightarrow B$ be a morphism of rings, and M a B -module. If M is flat over B and B is flat over A then M is flat over A .*

Proof. The functor $- \otimes_A M$ is isomorphic to the composition $(- \otimes_A B) \otimes_B M$ of exact functors. \square

Proposition 1.2.3. *Let $A \rightarrow B$ be a morphism of rings. If M is a flat A -module, then $B \otimes_A M$ is a flat B -module.*

Proof. The functor $- \otimes_B (B \otimes_A M)$ is isomorphic to the functor $- \otimes_A M$, which is exact. \square

¹See [3], chapter 2, section 2.7

Proposition 1.2.4. *Let $\varphi: A \rightarrow B$ be a morphism of rings, and M a B -module. M is flat over A if and only if for every $\mathfrak{q} \in \text{Specmax } B$ the module $M_{\mathfrak{q}}$ is flat over $A_{\mathfrak{p}}$, where $\mathfrak{p} = \varphi^{-1}\mathfrak{q}$.*

Proof. Notice that $\otimes_A M$ sends A -modules to B -modules, with the structure of B -module inherited from M . Let $\mathfrak{q} \in \text{Specmax } B$, and $\mathfrak{p} = \varphi^{-1}\mathfrak{q}$. We have an isomorphism of functors from the category of A -modules to the category of $B_{\mathfrak{q}}$ -modules:

$$\begin{aligned} (- \otimes_A M)_{\mathfrak{q}} &= (- \otimes_A M) \otimes_B B_{\mathfrak{q}} = - \otimes_A (M \otimes_B B_{\mathfrak{q}}) = \\ &= - \otimes_A M_{\mathfrak{q}} = - \otimes_A (A_{\mathfrak{p}} \otimes_{A_{\mathfrak{p}}} M_{\mathfrak{q}}) = (-)_{\mathfrak{p}} \otimes_{A_{\mathfrak{p}}} M_{\mathfrak{q}}. \end{aligned}$$

Localization is exact. Hence, if $\otimes_A M$ is flat then $\otimes_{A_{\mathfrak{p}}} M_{\mathfrak{q}}$ is exact. Conversely, if $\otimes_{A_{\mathfrak{p}}} M_{\mathfrak{q}}$ is exact for every $\mathfrak{q} \in \text{Specmax } B, \mathfrak{p} = \varphi^{-1}\mathfrak{q}$, then tensoring a short exact sequence $N_1 \rightarrow N_2 \rightarrow N_3$ with M we obtain a sequence of B -modules which is exact at every maximal ideal \mathfrak{q} . Therefore it is exact. \square

Proposition 1.2.5. *Let A be a ring. An A -module is flat if and only if $I \otimes_A M \rightarrow M$ is injective (equivalently, $\text{Tor}_1^A(A/I, M) = 0$) for every finitely generated ideal $I \subset A$.*

Proof. The “only if” part is trivial. We want to show that for arbitrary inclusion of A -modules $N' \subset N$ the induced morphism $N' \otimes_A M \rightarrow N \otimes_A M$ is injective.

We first show that $I \otimes_A M \rightarrow M$ is injective for every ideal I . Let $x \in I \otimes_A M$ be an element which vanishes in M . The element x is a finite linear combination of elementary tensors $y \otimes m$ where $y \in I, m \in M$. Thus there exists a finitely generated ideal $I' \subset I$ and $x' \in I' \otimes_A M$ such that the image of x' in $I \otimes_A M$ is equal to x . The map $I' \otimes_A M \rightarrow M$ is injective, so $x' = 0$ and hence $x = 0$, i.e. $I \otimes_A M \rightarrow M$ is injective. As a corollary, $\text{Tor}_1^A(N, M) = 0$ if N is a cyclic module, that is, $N = A/I$ for some ideal $I \subset A$.

Let N be an arbitrary module and N' its submodule. Consider an index set J whose elements are finite subsets of $N \setminus N'$. For $j \in J$ let N_j be the submodule of N generated by N' and j . If $j \subset j'$ then there is a natural injection $N_j \rightarrow N_{j'}$. The inclusion order on J makes it a directed poset. Clearly, $\text{colim}_{j \in J} N_j = N$.

Let $j \subset j'$ be an inclusion. Assume that $j' \setminus j$ consists of a single element. In this case $N_{j'}/N_j$ is a cyclic module. The short exact sequence $0 \rightarrow N_j \rightarrow N_{j'} \rightarrow N_{j'}/N_j \rightarrow 0$ induces an exact sequence $\text{Tor}_1^A(N_{j'}/N_j, M) \rightarrow N_j \otimes_A M \rightarrow N_{j'} \otimes_A M$. Since $N_{j'}/N_j$ is cyclic, $\text{Tor}_1^A(N_{j'}/N_j, M)$ vanishes, and so $N_j \otimes_A M \rightarrow N_{j'} \otimes_A M$ is injective.

A general inclusion $j \subset j'$ can be factored into a sequence of inclusions such that at each step only one new element appears. Hence $N_j \otimes_A M \rightarrow N_{j'} \otimes_A M$ is injective, which implies that the morphism $N' \otimes_A M \rightarrow \text{colim}_{j \in J} N_j \otimes_A M$ is injective too. It remains to recall that $\otimes_A M$ commutes with colimits. \square

Corollary 1.2.6. *Let A be a PID. An A -module M is flat if and only if it is torsion-free.*

Proposition 1.2.7. *Let A be a ring, let $0 \rightarrow M' \rightarrow M'' \rightarrow M \rightarrow 0$ be a short exact sequence of A -modules, and let N be an A -module. If M is flat then $M' \otimes_A N \rightarrow M'' \otimes_A N$ is injective.*

Proof. One can either refer to commutativity of Tor or do a direct proof as follows. Let $0 \rightarrow K \rightarrow F \rightarrow N \rightarrow 0$ be a short exact sequences with F a free module. Consider a commutative diagram with exact rows and columns:

$$\begin{array}{ccccccc}
 & & & & & & 0 \\
 & & & & & & \downarrow \\
 & & & & & & M' \otimes_A K \longrightarrow M'' \otimes_A K \longrightarrow M \otimes_A K \longrightarrow 0 \\
 & & & & & & \downarrow \\
 & & & & & & 0 \longrightarrow M' \otimes_A F \longrightarrow M'' \otimes_A F \longrightarrow M \otimes_A F \longrightarrow 0 \\
 & & & & & & \downarrow \\
 & & & & & & M' \otimes_A N \longrightarrow M'' \otimes_A N \longrightarrow M \otimes_A N \longrightarrow 0. \\
 & & & & & & \downarrow \\
 & & & & & & 0 \qquad \qquad \qquad 0 \qquad \qquad \qquad 0.
 \end{array}$$

A simple diagram chase finishes the proof. □

Theorem 1.2.8. *Let A be a local noetherian ring, and M an A -module of finite type. If M is flat then it is free.*

Proof. Let k be the residue field of A . Take a k -basis of $M \otimes_A k$. Lifting it to M we obtain a morphism from a free A -module F of finite type to M . By Nakayama lemma this morphism is surjective. Let K be its kernel. Tensoring the short exact sequence $0 \rightarrow K \rightarrow F \rightarrow M \rightarrow 0$ by k we obtain exact sequence $K \otimes_A k \rightarrow F \otimes_A k \rightarrow M \otimes_A k \rightarrow 0$. The morphism $K \otimes_A k \rightarrow F \otimes_A k$ is injective by proposition 1.2.7. The morphism $F \otimes_A k \rightarrow M \otimes_A k$ is an isomorphism by construction. Hence $K \otimes_A k$ is zero. On the other hand, K is of finite type since A is noetherian. So, Nakayama lemma shows that $K = 0$. □

1.3 Artin-Rees lemma and Krull intersection theorem

Let A be a ring, $I \subset A$ an ideal.

Definition 1.3.1. Let M be an A -module. An I -filtration on M is a descending chain of submodules $F_i M \subset M$, $i \in \mathbf{Z}_{\geq 0}$, such that $F_0 M = M$ and $IF_i M \subset F_{i+1} M$ for every i .

Definition 1.3.2. Let M be an A -module. An I -filtration $F_i M$ is called stable if $IF_i M = F_{i+1} M$ for sufficiently large i .

Proposition 1.3.3. Let A be a ring, $I \subset A$ an ideal, and let N, M be A -modules. If $F_i N$ is a stable I -filtration of N then the filtration of $N \otimes_A M$ by images of $F_i N \otimes_A M$ is stable.

Proof. Omitted. □

Proposition 1.3.4. Let $A \rightarrow B$ be a morphism of rings, $I \subset A$ an ideal, M a B -module, and $F_i M$ a stable I -filtration of M as an A -module. If each $F_i M$ is a B -submodule, then $F_i M$ is a stable IB -filtration of M as a B -module.

Proof. Omitted. □

Let M be an A -module endowed with an I -filtration $F_i M$. Consider a graded ring $B_I A = \bigoplus_{i=0}^{\infty} I^i$ and a $B_I A$ -module $B_F M = \bigoplus_{i=0}^{\infty} F_i M$.

Proposition 1.3.5. Let A be a noetherian ring, $I \subset A$ an ideal, M an A -module with an I -filtration $F_i M$. The filtration is stable if and only if $B_F M$ is of finite type over $B_I A$.

Lemma 1.3.6 (Artin-Rees lemma). Let A be a noetherian ring, $I \subset A$ an ideal, M an A -module with a stable I -filtration $F_i M$, and $N \subset M$ a submodule. The filtration $F_i N = N \cap F_i M$ is stable.

Proof. The ring $B_I A$ is noetherian since it is a quotient of the polynomial ring $A[x_1, \dots, x_n]$ for some n . The module $B_F N$ is a submodule of $B_F M$, and thus is of finite type. Now the claim follows from the previous proposition. □

Theorem 1.3.7 (Krull intersection theorem). Let A be a noetherian local ring, $I \subset A$ an ideal and M a module of finite type. If $F_i M$ is a stable I -filtration of M , then $\bigcap_{i=0}^{\infty} F_i M = 0$.

Proof. Consider the submodule $N = \bigcap_{i=0}^{\infty} F_i M$. By construction $N \cap F_i M = N$ for every i , and so by Artin-Rees lemma $N = IN$. Hence $N = \mathfrak{m}N$. Since N is of finite type, Nakayama lemma implies that $N = 0$. □

1.4 Modules of finite length

Let A be a ring, M a module. A strict chain of submodules of length n is an increasing sequence of submodules of M :

$$M_0 \subset M_1 \subset \dots \subset M_n,$$

such that $M_0 = 0$, $M_n = M$, and each inclusion $M_i \subset M_{i+1}$ is nontrivial.

We define $l_A(M)$, the length of M , as the supremum of lengths of strict chains.

Definition 1.4.1. M is called a module of finite length if $l_A(M)$ is finite (i.e. if the supremum exists).

Proposition 1.4.2. $l_A(M) = 1$ if and only if $M = A/\mathfrak{m}$ for some $\mathfrak{m} \in \text{Specmax } A$.

Proof. Excercise. □

Proposition 1.4.3. Let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be a short exact sequence of A -modules. If M is of finite length or M' and M'' are of finite length then all three modules are of finite length and $l_A(M) = l_A(M') + l_A(M'')$.

Proof. Excercise. □

Proposition 1.4.4. Let A be a ring, $\mathfrak{m} \subset A$ a maximal ideal of finite type, and M an A -module of finite type. If $\mathfrak{m}^n M = 0$ for some $n > 0$, then M is of finite length.

Proof. Let $n > 0$ be an integer. Suppose that A/\mathfrak{m}^n is of finite length. If a module M of finite type is annihilated by \mathfrak{m}^n then it is an A/\mathfrak{m}^n -module, and so is a quotient of a finite direct sum of A/\mathfrak{m}^n 's. Hence M is of finite length.

We next prove that A/\mathfrak{m}^n is of finite length using induction over n . The case $n = 1$ was already established. Consider a short exact sequence

$$0 \rightarrow \mathfrak{m}/\mathfrak{m}^n \rightarrow A/\mathfrak{m}^n \rightarrow A/\mathfrak{m} \rightarrow 0.$$

The module $\mathfrak{m}/\mathfrak{m}^n$ is of finite type since \mathfrak{m} is, and is annihilated by \mathfrak{m}^{n-1} , whence of finite length. But then A/\mathfrak{m}^n is also of finite length. □

Proposition 1.4.5. Let A be a ring, M an A -module. If $\text{Tor}_1^A(A/\mathfrak{m}, M) = 0$ for every $\mathfrak{m} \in \text{Specmax } A$, then $\text{Tor}_1^A(N, M) = 0$ for every module N of finite length.

Proof. We will do it by induction on $l_A(N)$. If $l_A(N) = 1$ then N is of the form A/\mathfrak{m} , and so $\text{Tor}_1^A(N, M) = 0$ by assumption. Otherwise there exists a proper nontrivial submodule $N' \subset N$. Consider an exact sequence $\text{Tor}_1^A(N', M) \rightarrow \text{Tor}_1^A(N, M) \rightarrow \text{Tor}_1^A(N/N', M)$ induced by short exact sequence $0 \rightarrow N' \rightarrow N \rightarrow N/N' \rightarrow 0$. Since $l_A(N') < l_A(N)$ and $l_A(N/N') < l_A(N)$, we see that $\text{Tor}_1^A(N', M) = \text{Tor}_1^A(N/N', M) = 0$, so $\text{Tor}_1^A(N, M) = 0$. □

1.5 Criteria of flatness

Theorem 1.5.1 (Critère local de platitude). Let $A \rightarrow B$ be a local morphism of noetherian local rings, k the residue field of A , and M a B -module of finite type. If $\text{Tor}_1^A(k, M) = 0$ then M is flat over A .

Proof. We want to show that for every ideal $I \subset A$ the module $\mathrm{Tor}_1^A(A/I, M)$ vanishes. Notice that if A/I is of finite length, then $\mathrm{Tor}_1^A(A/I, M) = 0$ by proposition 1.4.5.

Let $\mathfrak{m} \subset A$ be the maximal ideal, and $I \subset A$ an arbitrary ideal. Let $n > 0$ be an integer. Consider a diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & I \cap \mathfrak{m}^n & \longrightarrow & I & \longrightarrow & I/(I \cap \mathfrak{m}^n) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathfrak{m}^n & \longrightarrow & A & \longrightarrow & A/\mathfrak{m}^n \longrightarrow 0. \end{array}$$

Tensoring it with M over A we obtain a diagram

$$\begin{array}{ccccccc} (I \cap \mathfrak{m}^n) \otimes_A M & \longrightarrow & I \otimes_A M & \longrightarrow & (I/(I \cap \mathfrak{m}^n)) \otimes_A M \\ \downarrow & & \downarrow \alpha & & \downarrow \beta_n \\ \mathfrak{m}^n \otimes_A M & \longrightarrow & M & \longrightarrow & (A/\mathfrak{m}^n) \otimes_A M. \end{array}$$

with right exact rows. The cokernel of the map $I/(I \cap \mathfrak{m}^n) \rightarrow A/\mathfrak{m}^n$ is $A/(I + \mathfrak{m}^n)$. It has finite length by proposition 1.4.4. Thus $\mathrm{Tor}_1^A(A/(I + \mathfrak{m}^n), M) = 0$ and the morphism β_n is injective. As a consequence, $\ker(\alpha)$ is contained in the image of $(I \cap \mathfrak{m}^n) \otimes_A M$.

The filtration \mathfrak{m}^n on A is \mathfrak{m} -stable. Hence by Artin-Rees lemma the filtration $I \cap \mathfrak{m}^n$ on I is \mathfrak{m} -stable, and so the filtration on $I \otimes_A M$ by images of $(I \cap \mathfrak{m}^n) \otimes_A M$ is \mathfrak{m} -stable (notice that $I \otimes_A M$ is not necessarily an A -module of finite type!).

The module $I \otimes_A M$ has a structure of B -module via M , and the images of $(I \cap \mathfrak{m}^n) \otimes_A M$ in this module are B -submodules. Let $J = \mathfrak{m}B \subset B$. This ideal is proper since $A \rightarrow B$ is a local morphism. The filtration on $I \otimes_A M$ as a B -module is J -stable. Now, Krull intersection theorem tells us that $\ker(\alpha) = 0$ as a submodule of zero module. \square

Lemma 1.5.2. *Let $A \rightarrow B$ be a local morphism of local noetherian rings, $I \subset A$ an ideal, and M a B -module of finite type. If $\mathrm{Tor}_1^A(A/I, M) = 0$ and M/IM is a flat A/I -module, then M is a flat A -module.*

Proof. Let k be the residue field of A . A short exact sequence

$$0 \rightarrow K \rightarrow A/I \rightarrow k \rightarrow 0.$$

yields an exact sequence

$$\mathrm{Tor}_1^A(A/I, M) \rightarrow \mathrm{Tor}_1^A(k, M) \rightarrow K \otimes_A M \rightarrow A/I \otimes_A M$$

By assumptions $\mathrm{Tor}_1^A(A/I, M) = 0$. The modules K and A/I are A/I -modules, and the functor $\otimes_A M$ restricted to such modules is isomorphic to $\otimes_{A/I} M/IM$. The latter functor is exact, and so the arrow $K \otimes_A M \rightarrow A/I \otimes_A M$ is injective. Hence $\mathrm{Tor}_1^A(k, M) = 0$, and the local criterion of flatness finishes the proof. \square

Proposition 1.5.3. *Let A be a ring, M a flat A -module. If $M/\mathfrak{m}M \neq 0$ for every $\mathfrak{m} \in \text{Specmax } A$, then $N \otimes_A M = 0$ implies $N = 0$.*

Proof. If $\mathfrak{m} \in \text{Specmax } A$, then

$$(N \otimes_A M) \otimes_A k(\mathfrak{m}) = N/\mathfrak{m}N \otimes_{k(\mathfrak{m})} M/\mathfrak{m}M.$$

Since $N \otimes_A M = 0$, we see that $N/\mathfrak{m}N = 0$ for every $\mathfrak{m} \in \text{Specmax } A$. If N is of finite type, then by Nakayama $N_{\mathfrak{m}} = 0$ for every $\mathfrak{m} \in \text{Specmax } A$, so $N = 0$. If N is not of finite type, then we take an element $x \in N$ and consider a submodule N' generated by x . The morphism $N' \rightarrow N$ is injective, so $N' \otimes_A M \rightarrow N \otimes_A M$ is injective, and as a consequence $N' = 0$, i.e. $x = 0$. Hence, $N = 0$. \square

Theorem 1.5.4 (Critère de platitude par fibres, cas noethérien). *Let $A \rightarrow B \rightarrow C$ be local morphisms of local noetherian rings, and M a C -module of finite type. Let k be the residue field of A . If M is nonzero, flat over A , and $M \otimes_A k$ is flat over $B \otimes_A k$, then B is flat over A and M is flat over B .*

Proof. Let \mathfrak{m} be the maximal ideal of A , and $I = \mathfrak{m}B$. The natural map $\mathfrak{m} \otimes_A B \rightarrow I$ is surjective, and $(\mathfrak{m} \otimes_A B) \otimes_B C = \mathfrak{m} \otimes_A C$, so $\mathfrak{m} \otimes_A C \rightarrow I \otimes_B C$ is surjective. As a consequence, $\mathfrak{m} \otimes_A M \rightarrow I \otimes_B M$ is surjective.

The composition $\mathfrak{m} \otimes_A M \rightarrow I \otimes_B M \rightarrow M$ is injective, since M is flat over A . Hence $\mathfrak{m} \otimes_A M \rightarrow I \otimes_B M$ is an isomorphism, and $I \otimes_B M \rightarrow M$ is injective. In particular, $\text{Tor}_1^A(B/I, M) = 0$, so M is flat over B by lemma 1.5.2.

Consider an exact sequence $0 \rightarrow \mathfrak{m} \rightarrow A \rightarrow k \rightarrow 0$. Tensoring with B over A gives us an exact sequence $0 \rightarrow \text{Tor}_1^A(k, B) \rightarrow \mathfrak{m} \otimes_A B \rightarrow I \rightarrow 0$. Tensoring the latter sequence with M over B yields a sequence $0 \rightarrow \text{Tor}_1^A(k, B) \otimes_B M \rightarrow \mathfrak{m} \otimes_A M \rightarrow I \otimes_B M \rightarrow 0$. The last map is an isomorphism, so $\text{Tor}_1^A(k, B) \otimes_B M = 0$.

If $\mathfrak{m}_B \subset B$ and $\mathfrak{m}_C \subset C$ are maximal ideals, then $M/\mathfrak{m}_C M$ is nonzero by Nakayama, so $M/\mathfrak{m}_B M$ is nonzero. Hence, proposition 1.5.3 applies and shows that $\text{Tor}_1^A(k, B) = 0$. It remains to apply theorem 1.5.1. \square

1.6 Flatness in the context of schemes

Definition 1.6.1. Let $f: X \rightarrow Y$ be a morphism of schemes, and \mathcal{F} a sheaf of \mathcal{O}_X -modules. We say that \mathcal{F} is flat over Y at $x \in X$ if the stalk \mathcal{F}_x is a flat module over $\mathcal{O}_{Y, f(x)}$. We say that f is flat at $x \in X$ if \mathcal{O}_X is flat over Y at x . We say that \mathcal{F} is flat over Y if it is flat over Y at all points. We say that f is flat if \mathcal{O}_X is flat over Y .

Proposition 1.6.2. *Flat morphisms have following properties:*

(1) *If X and Y are affine schemes and \mathcal{F} is quasi-coherent, then \mathcal{F} is flat over Y if and only if $\Gamma(X, \mathcal{F})$ is a flat module over $\Gamma(Y, \mathcal{O}_Y)$.*

(2) Let $f: X \rightarrow Y$, $g: Y \rightarrow Z$ be morphisms, and \mathcal{F} a quasi-coherent sheaf. If \mathcal{F} is flat over Y and g is flat, then \mathcal{F} is flat over Z . In particular, a composition of flat morphisms is flat.

(3) Let $X \rightarrow Y$ be a morphism, \mathcal{F} a quasi-coherent sheaf, $g: Z \rightarrow Y$ a morphism, and $p: X \times_Y Z \rightarrow X$ a projection. If \mathcal{F} is flat over Y , then $p^*\mathcal{F}$ is flat over Z . In particular, a basechange of a flat morphism is flat.

(4) An open immersion is flat.

Proof. Follows easily from what we have already done. \square

Theorem 1.6.3. Let S, X, Y be locally noetherian schemes, and $f: X \rightarrow Y$ a morphism of schemes over S . Let \mathcal{F} a coherent \mathcal{O}_X -module. Assume that all stalks of \mathcal{F} are nonzero, \mathcal{F} is flat over S , and for every $s \in S$ the pullback of \mathcal{F} to X_s is flat over Y_s . Then \mathcal{F} is flat over Y and Y is flat over S at all points $y \in f(X)$.

Proof. Follows at once from theorem 1.5.4. \square

Corollary 1.6.4. Let S, X, Y be locally noetherian schemes. Let $f: X \rightarrow Y$ and $g: Y \rightarrow S$ be morphisms of schemes. If gf is flat and for every $s \in S$ the pullback $X_s \rightarrow Y_s$ of f is flat, then f is flat, and g is flat at all points $y \in f(X)$.

2 Étale morphisms

2.1 The module of Kähler differentials

Definition 2.1.1. Let $A \rightarrow B$ be a morphism of rings, and M a B -module. An A -derivation $d: B \rightarrow M$ is an A -module morphism, which satisfies Leibnitz identity: $d(b_1b_2) = b_2d(b_1) + b_1d(b_2)$ for every $b_1, b_2 \in B$.

Using Leibnitz identity we get $d(1) = d(1 \cdot 1) = d(1) + d(1)$, so that $d(1) = 0$. If $a \in A$, then $d(a) = ad(1) = 0$ by A -linearity.

A sum of two derivations is again an A -derivation, as well as a scalar multiple of a derivation by an element of B . Hence, A -derivations $B \rightarrow M$ form a B -module, which is denoted $\text{Der}_A(B, M)$. The association $M \mapsto \text{Der}_A(B, M)$ is a covariant functor in an evident way.

Theorem 2.1.2. Let $A \rightarrow B$ be a morphism of rings. The functor $\text{Der}_A(B, -)$ is representable.

Proof. Such proofs are better done on one's own. \square

The B -module which represents $\text{Der}_A(B, -)$ is denoted $\Omega_{B/A}^1$ and is called the module of Kähler differentials. The identity morphism $\text{id} \in \text{Hom}_B(\Omega_{B/A}^1, \Omega_{B/A}^1)$ gives rise to a derivation $d \in \text{Der}_A(B, \Omega_{B/A}^1)$ which is called the universal derivation. By construction, for every A -derivation $d' : B \rightarrow M$ there exists a unique B -module morphism $f : \Omega_{B/A}^1 \rightarrow M$ such that $d' = f \circ d$.

Proposition 2.1.3. *Let $A \rightarrow B \rightarrow C$ be morphisms of rings. There is an exact sequence of C -modules:*

$$\Omega_{B/A}^1 \otimes_B C \rightarrow \Omega_{C/A}^1 \rightarrow \Omega_{C/B}^1 \rightarrow 0.$$

Proof. Let M be a C -module. Consider a sequence

$$0 \rightarrow \text{Der}_B(C, M) \rightarrow \text{Der}_A(C, M) \rightarrow \text{Der}_A(B, M).$$

The first map takes a B -derivation $d : C \rightarrow M$ and views it as an A -derivation. The second map precomposes an A -derivation $d : C \rightarrow M$ with $B \rightarrow C$. Clearly, the first map is injective. On the other hand, if an A -derivation $d : C \rightarrow M$ vanishes when restricted to B , then it is a B -derivation. Hence, the sequence is exact for every module M . Since it is also natural in M , we obtain an exact sequence of functors:

$$0 \rightarrow \text{Hom}_C(\Omega_{C/B}^1, -) \rightarrow \text{Hom}_C(\Omega_{C/A}^1, -) \rightarrow \text{Hom}_B(\Omega_{B/A}^1, -).$$

But $\text{Hom}_B(\Omega_{B/A}^1, -) = \text{Hom}_C(\Omega_{B/A}^1 \otimes_B C, -)$, and so this sequence of functors yields the desired sequence of modules. \square

Proposition 2.1.4. *Let $A \rightarrow B$ be a morphism of rings, $I \subset B$ an ideal. There is an exact sequence of B/I -modules:*

$$I/I^2 \rightarrow \Omega_{B/A}^1 \otimes_B B/I \rightarrow \Omega_{(B/I)/A}^1 \rightarrow 0.$$

Proof. Let M be a B/I -module. Consider a sequence:

$$0 \rightarrow \text{Der}_A(B/I, M) \rightarrow \text{Der}_A(B, M) \rightarrow \text{Hom}_{B/I}(I/I^2, M)$$

Here the first map takes a derivation $B/I \rightarrow M$ and precomposes it with the quotient morphism $B \rightarrow C$. The second map takes a derivation $d : B \rightarrow M$ and restricts it to I . Leibnitz identity and the fact that $IM = 0$ show that d vanishes on I^2 , and determines a morphism $d : I/I^2 \rightarrow M$ of B/I -modules. On the other hand, if $d|_I = 0$, then clearly d comes from an A -derivation $B/I \rightarrow M$, hence the sequence is exact. We conclude as in the previous proposition. \square

Proposition 2.1.5. *Let $A \rightarrow B$ be a morphism of rings. If it is of finite type, then $\Omega_{B/A}^1$ is of finite type over B .*

Proof. Let $B = A[x]$, the free polynomial algebra in one variable. Due to Leibnitz identity, an A -derivation $d: B \rightarrow M$ is determined by $d(x)$. Direct check shows that any value $d(x) \in M$ is admissible. Hence, the functor $\text{Der}_A(B, -)$ is represented by B .

Now proposition 2.1.3 immediately implies that if B is a free polynomial A -algebra in finitely many variables, then $\Omega_{B/A}^1$ is of finite type. Then proposition 2.1.4 shows that the same holds when B is a quotient of such an algebra. \square

Proposition 2.1.6. *Let $A \rightarrow B$ be a morphism of rings, $s \in B$ a unit, $b \in B$ an element, and $d: B \rightarrow M$ an A -derivation.*

$$d\left(\frac{b}{s}\right) = \frac{sdb - bds}{s^2}$$

Proof. From the formula $0 = d(1) = d(ss^{-1}) = sd(s^{-1}) + s^{-1}ds$ we conclude that $d\left(\frac{1}{s}\right) = -\frac{ds}{s^2}$, and then the claim follows by Leibnitz identity. \square

Proposition 2.1.7. *Let A be a ring, $S \subset A$ a multiplicative system, and $A \rightarrow A_S$ a localization morphism. $\Omega_{A_S/A}^1 = 0$.*

Proof. From the previous proposition it follows that every A -derivation of A_S is zero. \square

Proposition 2.1.8. *Let $A \rightarrow B$ be a morphism of rings, and $S \subset B$ a multiplicative system. The natural morphism $\Omega_{B/A}^1 \otimes_B B_S \rightarrow \Omega_{B_S/A}^1$ is an isomorphism.*

Proof. Let M be a B_S -module. Consider the morphism $\text{Der}_A(B_S, M) \rightarrow \text{Der}_A(B, M)$ induced by the ring morphism $B \rightarrow B_S$. We want to show that it is surjective. Let $d: B \rightarrow M$ be an A -derivation. It induces a derivation $D: B_S \rightarrow M$ by the rule

$$D\left(\frac{b}{s}\right) = \frac{sdb - bds}{s^2}.$$

Additivity and Leibnitz identity follow from trivial but lengthy calculations. Clearly, $D\left(\frac{b}{1}\right) = \frac{db}{1}$, so D is an A -derivation which restricts to d on B .

Hence the restriction morphism $\text{Der}_A(B_S, -) \rightarrow \text{Der}_A(B, -)$ is surjective, and we obtain an exact sequence of functors

$$0 \rightarrow \text{Der}_B(B_S, -) \rightarrow \text{Der}_A(B_S, -) \rightarrow \text{Der}_A(B, -) \rightarrow 0.$$

The observation that $\text{Der}_B(B_S, -) = 0$ finishes the proof. \square

Proposition 2.1.9. *Let A be a ring, let B, C be A -algebras, and let $D = B \otimes_A C$. The morphism $\Omega_{B/A}^1 \otimes_B D \rightarrow \Omega_{B \otimes_A C/C}^1$ induced by ring morphism $B \rightarrow D$ is an isomorphism. In other words, Ω^1 is stable under base change.*

Proof. Let M be a module over D . Consider a morphism $\text{Der}_C(D, M) \rightarrow \text{Der}_A(B, M)$ induced by ring morphism $B \rightarrow D$. An element of $\text{Der}_C(D, M)$ is a bilinear map $d: B \times C \rightarrow M$ which satisfies the following identities for every $a \in A, b \in B, c \in C, b_i \in B$:

$$\begin{aligned} d(ab, c) &= d(b, ac) = ad(b, c), \\ d(b, c) &= (1 \otimes_A c)d(b, 1), \\ d(b_1 b_2, 1) &= (b_1 \otimes_A 1)d(b_2, 1) + (b_2 \otimes_A 1)d(b_1, 1). \end{aligned}$$

From this description it is clear that if d vanishes in $\text{Der}_A(B, M)$, then $d = 0$. Given $D \in \text{Der}_A(B, M)$ we define $d(b, c) = (1 \otimes_A c)D(b)$, which clearly satisfies the equation above. Hence $\text{Der}_C(D, M) \rightarrow \text{Der}_A(B, M)$ is an isomorphism for every M . \square

Proposition 2.1.10. *Let $f: A \rightarrow B$ be a morphism of rings, $S \subset B$ a multiplicative system. The natural morphism $\Omega_{B/A}^1 \otimes_B B_S \rightarrow \Omega_{B_S/A_{f^{-1}S}}^1$ induced by ring morphisms $B \rightarrow B_S$ and $A \rightarrow A_{f^{-1}S}$ is an isomorphism.*

Proof. The natural morphism in question factors as $\Omega_{B/A}^1 \otimes_B B_S \rightarrow \Omega_{B_S/A}^1 \rightarrow \Omega_{B_S/A_{f^{-1}S}}^1$. The first morphism is an isomorphism as proposition 2.1.8 shows. On the other hand, $B_S \otimes_A A_{f^{-1}S} = B_S$, so that the second morphism is an isomorphism by proposition 2.1.9. \square

Let $f: X \rightarrow Y$ be a morphism of schemes. One can extend the definition of Ω^1 to $X \rightarrow Y$ in two ways. First, since $\Omega_{B/A}^1$ commutes with restrictions to principal open subsets of $\text{Spec } B$ and pullbacks to principal open subsets of $\text{Spec } A$, one can pick a covering U_i of Y by open affines and coverings V_{ij} of $f^{-1}U_i$ by open affines, then glue various Ω_{V_{ij}/U_i}^1 , and show that this construction does not depend on the choice of covers. The other way is, given a morphism $f: (X, \mathcal{O}_X) \rightarrow (Y, \mathcal{O}_Y)$ of ringed spaces and a \mathcal{O}_X -module \mathcal{F} , define a \mathcal{O}_X -module of derivations $\text{Der}_{f^{-1}\mathcal{O}_Y}(\mathcal{O}_X, \mathcal{F})$. One then shows that whenever X, Y are schemes and the morphism f is local, $\text{Der}_{f^{-1}\mathcal{O}_Y}(\mathcal{O}_X, -)$ is represented by a quasi-coherent \mathcal{O}_X -module, which agrees with Ω^1 when X and Y are affine. Either way, one obtains the following theorem:

Theorem 2.1.11. *To every morphism of schemes $f: X \rightarrow Y$ one can associate a quasi-coherent \mathcal{O}_X -module $\Omega_{X/Y}^1$ which has following properties:*

- *If X, Y are affine, then $\Omega_{X/Y}^1$ coincides with the module of Kähler differentials associated to the ring morphism $\Gamma(Y, \mathcal{O}_Y) \rightarrow \Gamma(X, \mathcal{O}_X)$.*
- *$\Omega_{X/Y}^1$ commutes with restrictions to opens $U \subset X$.*
- *Let $X \xrightarrow{f} S$ and $Y \xrightarrow{g} S$ be morphism. The sheaf $\Omega_{X \times_S Y/Y}^1$ is isomorphic to $p^* \Omega_{X/S}^1$, where $p: X \times_S Y \rightarrow X$ is a projection.*

- If $X \xrightarrow{f} Y \xrightarrow{g} Z$ are morphisms, then there is an exact sequence

$$f^*\Omega_{Y/Z}^1 \rightarrow \Omega_{X/Z}^1 \rightarrow \Omega_{X/Y}^1 \rightarrow 0.$$

- If $X \xrightarrow{f} Y$ is a morphism and $Z \xrightarrow{g} X$ is a closed immersion with ideal sheaf \mathcal{I} , then there exists an exact sequence

$$\mathcal{I}/\mathcal{I}^2 \rightarrow g^*\Omega_{X/Y}^1 \rightarrow \Omega_{Z/Y}^1 \rightarrow 0.$$

- If $f: X \rightarrow Y$ is locally of finite type, then $\Omega_{X/Y}^1$ is locally of finite type (in particular, coherent if X is locally noetherian).

2.2 Étale algebras over fields

Proposition 2.2.1. *Let $k \rightarrow K$ be a finite extension of fields. $\Omega_{K/k}^1$ vanishes if and only if $k \rightarrow K$ is separable.*

Proof. Assume that $k \rightarrow K$ is finite and separable. Let $x \in K$ be a primitive element, f its minimal polynomial. Let M be a K -module, and $d: K \rightarrow M$ a derivation.

$$0 = d(f(x)) = f'(x)dx.$$

Since K is separable, $f'(x) \neq 0$, so $dx = 0$ in M . Since K is generated over k by powers of x , we conclude that $d = 0$.

Assume that $k \rightarrow K$ is inseparable and primitive. Let $x \in K$ be a primitive element and f its minimal polynomial. Write $K = k[T]/(f)$. Recall that every derivation $d \in \text{Der}_k(k[T], K)$ is determined by $d(T)$ and $d(T)$ can be arbitrary. Set $d(T) = x$. Then d vanishes when restricted to (f) , since $d(gf) = g(x)f'(x)dx + f(x)dg = 0$ as $f(x) = 0$ and $f'(x) = 0$. Hence d comes from some derivation in $\text{Der}_k(k[T]/(f), K)$ i.e. $\text{Der}_k(K, K)$. As a consequence, $\text{Der}_k(K, K) \neq 0$.

Assume that $k \rightarrow K$ is inseparable. There is a nontrivial proper subfield $E \subset K$ such that $E \rightarrow k$ is inseparable and primitive. Then $\Omega_{K/k}^1$ is nonzero, since its quotient $\Omega_{K/E}^1$ is nonzero. \square

Proposition 2.2.2. *Let k be an algebraically closed field, A a k -algebra of finite type, and $\mathfrak{m} \in \text{Specmax } A$. The homomorphism $\mathfrak{m}/\mathfrak{m}^2 \rightarrow \Omega_{A/k}^1 \otimes_A k(\mathfrak{m})$ is an isomorphism.*

Proof. We need to prove that the natural restriction map

$$\text{Der}_k(A, M) \rightarrow \text{Hom}_{A/\mathfrak{m}}(\mathfrak{m}/\mathfrak{m}^2, M)$$

is an isomorphism for every A/\mathfrak{m} -module M .

By Hilbert's Nullstellensatz the composition $k \rightarrow A \rightarrow A/\mathfrak{m}$ is an isomorphism. In particular, $\text{Der}_k(A/\mathfrak{m}, -) = 0$, so that the natural map in question is injective. Let $f: \mathfrak{m}/\mathfrak{m}^2 \rightarrow M$ be a morphism of A/\mathfrak{m} -modules. We define a map $d: A \rightarrow M$ by sending an element $a \in A$ to $f(a - a(\mathfrak{m}))$, where $a(\mathfrak{m})$ is the image of a modulo \mathfrak{m} interpreted as an element of A . If $a_1, a_2 \in A$, then

$$\begin{aligned} a_1 a_2 - a_1(\mathfrak{m})a_2(\mathfrak{m}) &= \\ &= (a_1 - a_1(\mathfrak{m}))(a_2 - a_2(\mathfrak{m})) + a_2(\mathfrak{m})(a_1 - a_1(\mathfrak{m})) + a_1(\mathfrak{m})(a_2 - a_2(\mathfrak{m})). \end{aligned}$$

Also, $a_i = a_i(\mathfrak{m})$ in A/\mathfrak{m} , so that $d(a_1 a_2) = a_2 d(a_1) + a_1 d(a_2)$. Clearly, d vanishes on elements of k , so it is a derivation. \square

Definition 2.2.3. Let k be a field. A k -algebra A is called étale if it is a finite cartesian product of finite separable extensions of k .

Theorem 2.2.4. Let k be a field. A k -algebra of finite type A is étale if and only if $\Omega_{A/k}^1 = 0$.

Proof. Let A be a k -algebra of finite type such that $\Omega_{A/k}^1 = 0$. Let us first assume that k is algebraically closed. By virtue of proposition 2.2.2 we then know that $\mathfrak{m}/\mathfrak{m}^2 = 0$ for every maximal ideal \mathfrak{m} of A . Localizing at \mathfrak{m} and applying Nakayama lemma we conclude that $A_{\mathfrak{m}}$ is a field, the kernel of the localization morphism $A \rightarrow A_{\mathfrak{m}}$ is \mathfrak{m} , and $A_{\mathfrak{m}} = A/\mathfrak{m}$. By Nullstellensatz, $A/\mathfrak{m} \cong k$.

Let $\mathfrak{p} \in \text{Spec } A$ be a prime, and let \mathfrak{m} be a maximal ideal containing it. Let $a \in \mathfrak{m}$. Since a vanishes in $A_{\mathfrak{m}}$, there exists $s \notin \mathfrak{m}$ such that $sa = 0$ in A . In particular, $sa \in \mathfrak{p}$, so $a \in \mathfrak{p}$. Hence each prime of A is maximal.

The algebra A is noetherian, so that the set of its minimal primes is finite. But all primes are maximal, so $\text{Specmax } A$ is finite. Now, consider a morphism

$$A \rightarrow \prod_{\mathfrak{m} \in \text{Specmax } A} A/\mathfrak{m} \quad (1)$$

By Chinese remainder theorem it is surjective. But $A/\mathfrak{m} = A_{\mathfrak{m}}$, so that the kernel of this morphism consists of elements which vanish in all localizations of A at maximal ideals, i.e. the kernel is zero. Hence, this morphism is an isomorphism. In particular, $\dim_k A$ is finite.

Now, let k be arbitrary, and \bar{k} its algebraic closure. Let $A_{\bar{k}} = A \otimes_k \bar{k}$. Since $\dim_{\bar{k}} A_{\bar{k}}$ is finite, $\dim_k A$ is finite too. Let $\mathfrak{p} \in \text{Spec } A$ be a prime. The k -algebra A/\mathfrak{p} is finite-dimensional and has no zero divisors, hence it is a field. So $\text{Spec } A = \text{Specmax } A$, and $\text{Specmax } A$ is finite.

We consider a morphism as in (1). Its kernel is the nilradical of A . If $a \in A$ is nilpotent, then its image in $A_{\bar{k}}$ is nilpotent too, hence zero. But $A \rightarrow A_{\bar{k}}$ is injective, so that the kernel of (1) is zero. Now, proposition 2.2.1 finishes the proof. \square

2.3 Unramified morphisms

Definition 2.3.1. Let $f: X \rightarrow Y$ be a morphism of schemes. We say that f is unramified if f is locally of finite type and $\Omega_{X/Y}^1 = 0$.

Proposition 2.3.2. *Unramified morphisms have following properties:*

- (1) *If $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ are unramified, then gf is unramified.*
- (2) *If $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ are such that gf is unramified, then f is unramified.*
- (3) *If $f: X \rightarrow S$ is unramified, and $g: Y \rightarrow S$ is a morphism, then the pullback $X \times_S Y \rightarrow Y$ of f is unramified.*
- (4) *Open immersions are unramified.*

Proof. (1) The composition gf is locally of finite type. The exact sequence

$$g^* \Omega_{Y/Z}^1 \rightarrow \Omega_{X/Z}^1 \rightarrow \Omega_{X/Y}^1 \rightarrow 0$$

implies that $\Omega_{X/Z}^1 = 0$.

(2) The exact sequence above shows that $\Omega_{X/Y}^1 = 0$. The fact that f is locally of finite type is left as an exercise (see [2] tag 01T8).

(3) Follows from proposition 2.1.9.

(4) Follows from proposition 2.1.7. □

Proposition 2.3.3. *Let $f: X \rightarrow Y$ be a morphism locally of finite type. It is unramified if and only if for each $y \in Y$ the fiber $X_y \rightarrow y$ is unramified.*

Proof. If $\Omega_{X/Y}^1 = 0$, then clearly each fiber is unramified. Conversely, if $X_y \rightarrow y$ is unramified, then the fiber of $\Omega_{X/Y}^1$ at each point $x \in X$ is zero, as an inclusion of a point $x \in X$ factors through $X_{f(y)} \rightarrow X$. Since $\Omega_{X/Y}^1$ is locally of finite type, Nakayama lemma shows that $\Omega_{X/Y}^1 = 0$. □

Proposition 2.3.4. *Let $X \rightarrow \text{Spec } k$ be a scheme over a field. It is unramified if and only if X is discrete as a topological space, and for every $x \in X$ the field extension $k \rightarrow k(x)$ is finite separable.*

Proof. Assume that $X \rightarrow \text{Spec } k$ is unramified. Let $x \in X$ and $U \subset X$ be an affine open neighbourhood of x which is of finite type over $\text{Spec } k$. By theorem 2.2.4 we conclude that U is a spectrum of an étale algebra over k . In particular, U is discrete. Hence X is discrete.

Assuming the converse, take $x \in X$ and $U \subset X$ an affine open neighbourhood of x . Since X is discrete, U is discrete too, and as U is quasi-compact, we conclude that U is finite as a topological space. Hence U is a spectrum of an étale algebra over k , and so $\Omega_{X/k}^1|_U = 0$. As a consequence, $\Omega_{X/k}^1 = 0$. Since U is a spectrum of an algebra of finite type over k , we conclude that $X \rightarrow \text{Spec } k$ is locally of finite type. \square

Proposition 2.3.5. *Let X, Y be schemes and $f: X \rightarrow Y$ a morphism locally of finite type. The fiber of $\Omega_{X/Y}^1$ at x is zero if and only if the residue field extension $k(f(x)) \rightarrow k(x)$ is finite separable, and $\mathfrak{m}_{Y, f(y)} \mathcal{O}_{X, x} = \mathfrak{m}_{X, x}$.*

Proof. We immediately reduce to the case when $X = \text{Spec } B$ and $Y = \text{Spec } A$ are affine, and f is of finite type. Let $\mathfrak{q} \in \text{Spec } B$ and $\mathfrak{p} = f(\mathfrak{q})$.

Assume that $\Omega_{B/A}^1 \otimes_B k(\mathfrak{q}) = 0$. Since $\Omega_{B/A}^1$ is of finite type, Nakayama lemma implies that $(\Omega_{B/A}^1)_{\mathfrak{q}} = 0$. Hence replacing B by its localization at some element not contained in \mathfrak{q} we may assume that $\Omega_{B/A}^1 = 0$. As a consequence, $\Omega_{B_{\mathfrak{p}}/A_{\mathfrak{p}}}^1 = 0$.

Consider a ring $B \otimes_A k(\mathfrak{p})$. Since $\Omega_{B \otimes_A k(\mathfrak{p})/k(\mathfrak{p})}^1 = 0$ and B is of finite type over A , theorem 2.2.4 shows that $B \otimes_A k(\mathfrak{p})$ is a finite étale algebra over $k(\mathfrak{p})$.

The morphism $A_{\mathfrak{p}} \rightarrow k(\mathfrak{p})$ is surjective, so $B_{\mathfrak{q}} \rightarrow B_{\mathfrak{q}} \otimes_{A_{\mathfrak{p}}} k(\mathfrak{p})$ is surjective. On the other hand

$$B \otimes_A k(\mathfrak{p}) = B \otimes_A (A_{\mathfrak{p}} \otimes_{A_{\mathfrak{p}}} k(\mathfrak{p})) = B_{\mathfrak{p}} \otimes_{A_{\mathfrak{p}}} k(\mathfrak{p}),$$

so $B_{\mathfrak{q}} \otimes_{A_{\mathfrak{p}}} k(\mathfrak{p})$ is a localization of a finite étale algebra over $k(\mathfrak{p})$, hence is itself such an algebra.

The morphism $\text{Spec}(B_{\mathfrak{q}} \otimes_{A_{\mathfrak{p}}} k(\mathfrak{p})) \rightarrow \text{Spec } B_{\mathfrak{q}}$ is a closed immersion. In particular, it is injective and sends closed points to closed points. As $B_{\mathfrak{q}}$ has only one maximal ideal, we conclude that $B_{\mathfrak{q}} \otimes_{A_{\mathfrak{p}}} k(\mathfrak{p})$ also has unique maximal ideal, which forces it to be a finite separable field extension of $k(\mathfrak{p})$. On the other hand $B_{\mathfrak{q}} \otimes_{A_{\mathfrak{p}}} k(\mathfrak{p}) = B_{\mathfrak{q}}/\mathfrak{p}B_{\mathfrak{q}}$, so that $\mathfrak{p}B_{\mathfrak{q}} = \mathfrak{q}B_{\mathfrak{q}}$.

Now, assume that $\mathfrak{p}B_{\mathfrak{q}} = \mathfrak{q}B_{\mathfrak{q}}$ is maximal, and that $k(\mathfrak{q})$ is a finite separable extension of $k(\mathfrak{p})$. Our assumptions imply that $B_{\mathfrak{q}}/\mathfrak{p}B_{\mathfrak{q}} = B_{\mathfrak{q}} \otimes_{A_{\mathfrak{p}}} k(\mathfrak{p}) = k(\mathfrak{q})$. Hence $\Omega_{B/A}^1 \otimes_B k(\mathfrak{q}) = \Omega_{B_{\mathfrak{q}}/A_{\mathfrak{p}}}^1 \otimes_{B_{\mathfrak{q}}} k(\mathfrak{q}) = \Omega_{k(\mathfrak{q})/k(\mathfrak{p})}^1 = 0$. \square

2.4 Étale morphisms

Definition 2.4.1. Let $f: X \rightarrow Y$ be a morphism of schemes. We say that f is étale if it is unramified and flat.

Proposition 2.4.2. *Étale morphisms have following properties:*

- (1) *If $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ are étale, then gf is étale.*
- (2) *If $f: X \rightarrow S$ is étale, and $g: Y \rightarrow S$ is a morphism, then the pullback $X \times_S Y \rightarrow Y$ of f is étale.*
- (3) *Open immersions are étale.*
- (4) *If a morphism $f: X \rightarrow Y$ of schemes is locally of finite type, flat, and every fiber $X_y \rightarrow y$ is unramified, then f is étale.*

Proof. Everything follows at once from corresponding properties of flat and unramified morphisms. \square

Proposition 2.4.3. *Let $f: X \rightarrow Y$ and $g: Y \rightarrow S$ be morphisms of schemes. If gf is étale and g is unramified, then f is étale. If in addition f is surjective, then g is étale.*

Proof. Follows from corollary 1.6.4 because each fiber Y_s is a disjoint union of spectra of fields. \square

References

- [1] D. Eisenbud, *Commutative Algebra with a View Toward Algebraic Geometry*, Graduate Texts in Mathematics **150**, Springer, New York, 1995
- [2] A.J. de Jong et al., *Stacks Project*
- [3] C. Weibel, *An Introduction to Homological Algebra*, Cambridge University Press, Cambridge, 1994